

Permutations as Minimal Powers of a Single-cycle Class-sum

JACOB KATRIEL

The structure-constant $[(p)]_n \cdot [(1)^{i_1}(2)^{i_2} \cdots (n)^{i_n}]_n$ that corresponds to the merging of the p cycles $(1)^{k_1}(2)^{k_2} \cdots (u-1)^{k_{u-1}}$, $(\sum_{i=1}^{u-1} k_i = p)$ into a single cycle of length $u = \sum_{i=1}^{u-1} i k_i$ is derived. The result is used to evaluate the structure constant $[(p)]_n^k$ where k is the minimal power of $[(p)]_n$ generating the class-sum $[(1)^{i_1}(p)^{i_2} \cdots ((i-1)(p-1)+1)^{i_n}]_n$ (i.e. $\sum_{i=1}^n (i-1)i_i = k$).

© 1995 Academic Press Limited

1. INTRODUCTION

The class-algebra of the symmetric group has attracted considerable attention because of its representation-theoretic applications, as well as those in combinatorics, graph theory, the quantum-mechanical many-body problem and in other areas. The product of a pair of class-sums C_1 and C_2 is a linear combination with non-negative integral coefficients $C_1 \cdot C_2|_{C_i}$ of class-sums C_i , which we write in the form

$$C_1 \cdot C_2 = \sum_{C_i \in CS_n} C_i \cdot C_2|_{C_i} C_i$$

These coefficients are sometimes referred to as the structure constants of the class-algebra. Several attempts have been reported to derive these coefficients using their combinatorial significance [2–6, 11]. Closely related work involving the formulation of covering theorems, which specify the non-vanishing structure constants without explicitly evaluating them, was presented by Arad and Herzog [1].

A combinatorial study of the product of the class of transpositions with an arbitrary class of the symmetric group [10], followed by a similar study of the class of three-cycles [8], motivated the formulation of a conjecture concerning the form of the product of a single-cycle class-sum of arbitrary length with an arbitrary class-sum in the symmetric group algebra [9].

In the present paper, the structure-constant

$$[(p)]_n \cdot [(1)^{i_1}(2)^{i_2} \cdots (n)^{i_n}]_n$$

that corresponds to the merging of the p cycles $(1)^{k_1}(2)^{k_2} \cdots (u-1)^{k_{u-1}}$, $(\sum_{i=1}^{u-1} k_i = p)$ into a single cycle of length $u = \sum_{i=1}^{u-1} i k_i$ is derived. The result is used to evaluate the structure constant $[(p)]_n^k$ where k is the minimal power of $[(p)]_n$ generating the class-sum $[(1)^{i_1}(p)^{i_2} \cdots ((i-1)(p-1)+1)^{i_n}]_n$ (i.e. $\sum_{i=1}^n (i-1)i_i = k$). This is a generalization of the case $p = 2$ that was studied by Dénes [6], the graph-theoretical ramifications of which were recently elaborated upon by Moszkowski [12] and by Goulden and Pepper [7].

2. ON PRODUCTS WITH A MINIMAL NUMBER OF CYCLES OF A SINGLE-CYCLE WITH AN ARBITRARY CLASS-SUM

When a permutation of type $\pi^* \equiv (1)^{i_1}(2)^{i_2} \cdots (n)^{i_n}$ is multiplied by a p -cycle which possesses one index in common with each one of p cycles in π^* , it is found that these p cycles merge into one, i.e.,

$$(i_1, i_2, \dots, i_p) \cdot \underbrace{(i_1, \dots)}_{\lambda_1} \underbrace{(i_2, \dots)}_{\lambda_2} \cdots \underbrace{(i_p, \dots)}_{\lambda_p} * * * = \overbrace{(i_1, \dots, i_2, \dots, \dots, i_p, \dots)}^u * * *$$

where *** denotes the cycles that have no index in common with the p -cycle (i_1, i_2, \dots, i_p) and the symbol

$$\underbrace{i_j, \dots}_{\lambda_j}$$

denotes a specific sequence of λ_j indices, beginning with i_j . $\Lambda = \{\lambda_i; i = 1, 2, \dots, p\}$ is the p -tuple of lengths of the p cycles that are being merged. The length of the cycle generated is $u = \sum_{i=1}^p \lambda_i = \sum_{i=1}^{u-1} i \cdot k_i$, where k_i is the number of cycles of length i in the set Λ . Π denotes the particular sequence of u integers

$$\underbrace{i_1, \dots}_{\lambda_1}, \underbrace{i_2, \dots}_{\lambda_2}, \dots, \underbrace{i_p, \dots}_{\lambda_p}$$

The resulting permutation is of type

$$\pi = (1)^{l_1-k_1}(2)^{l_2-k_2} \dots (u-1)^{l_{u-1}-k_{u-1}}(u)^{l_u+1} \dots (n)^{l_n}.$$

The class-sum of permutations of type π will be denoted by $[\pi]_n$. We now prove the following.

THEOREM 1.

$$[(p)]_n \cdot [(1)^{l_1}(2)^{l_2} \dots (n)^{l_n}]_n \Big|_{[(1)^{l_1-k_1}(2)^{l_2-k_2} \dots (u)^{l_u+1} \dots (n)^{l_n}]_n} = u \frac{(p-1)!}{\prod_{i=1}^{u-1} k_i!} (l_u + 1)$$

where $\sum_{i=1}^n il_i = n$, $\sum_{i=1}^{u-1} ik_i = u$ and $\sum_{i=1}^{u-1} k_i = p$.

PROOF. In order to count the number of ways in which the particular cycle (Π) could be obtained as a product of some p -cycle with an appropriate permutation of type π , we note that i_1 could be selected within Π in u different ways. Independently, we can arrange the cycle lengths $\lambda_1, \lambda_2, \dots, \lambda_p$ in $p!/\prod_{i=1}^{u-1} k_i!$ distinct ways. Having selected the initial index i_1 and a particular ordering of the cycle lengths we obtain a well-defined set of cycles by appropriately splitting (Π) , as well as a set of indices comprising the p -cycle (i_1, i_2, \dots, i_p) . This p -cycle arises in p equivalent forms, related by cyclic permutations, by making all the different choices for i_1 . To eliminate this multiple counting we divide by p .

Finally, assuming that the original permutation had l_u u -cycles, each one of the $l_u + 1$ u -cycles in the permutation π can be obtained from appropriate Λ -sets of cycles multiplied by appropriate p -cycles. Since these $l_u + 1$ u -cycles consist of disjoint sets of indices, they give rise to $l_u + 1$ distinct sets of products of p -cycles by permutations of type π^* .

Multiplying the factors u , $p!/\prod_{i=1}^{u-1} k_i!$, $1/p$ and $l_u + 1$ specified above, the theorem follows. \square

We note in passing that this theorem is an immediate consequence of the conjecture presented in [9] concerning products of a single-cycle by arbitrary class-sums in the algebra of the symmetric group.

3. COEFFICIENTS OF CLASS-SUMS WITH A MINIMAL NUMBER OF CYCLES, IN THE EXPANSION OF $[(p)]_n^k$

In this section we evaluate the structure constants

$$[(p)]_n^k \Big|_{[(1)^{l_1}(2)^{l_2} \dots (n)^{l_n}]_n},$$

where $[(1)^1(2)^2 \cdots (n)^n]_n$ is a class with the smallest possible number of cycles within the sets of classes in the expansion of $[(p)]_n^k$. Since the number of cycles can be reduced by $p-1$ upon each application of $[(p)]_n$ (i.e. by merging p cycles into one), the minimal number of cycles in $[(p)]_n^k$ is obtained if such merging takes place k times, starting from the identity, $[(1)^n]_n$, which consists of n cycles. Thus, the minimal number of cycles in class-sums appearing in the expansion of $[(p)]_n^k$ is $n - (p-1)k$.

Repeated fusions of p cycles at a time, starting from $[(p)]_n$, can only give rise to cycles the lengths of which can be written in the form $i(p-1)+1$, where i is a non-negative integer. The class-sums appearing after k such fusions, i.e. in $[(p)]_n^k$, will be specified in terms of the indices $l_1, l_2, \dots, l_{i_{\max}}$, where l_i is the number of cycles of length $(i-1)(p-1)+1$, and

$$i_{\max} = 1 + \min\left(\left\lfloor \frac{n-1}{p-1} \right\rfloor, k\right).$$

The total number of cycles satisfies $\sum_{i=1}^{i_{\max}} l_i = n - k(p-1)$, which, noting that $\sum_{i=1}^{i_{\max}} ((i-1)(p-1)+1)l_i = n$, can be written in the more convenient form

$$\sum_{i=1}^{i_{\max}} (i-1)l_i = k. \quad (1)$$

We shall now prove the following.

THEOREM 2.

$$[(p)]_n^k \Big| \prod_{j=1}^{i_{\max}} ((j-1)(p-1)+1)^{l_j} \Big|_n = k! \prod_{j=2}^{i_{\max}} \left(\frac{((j-1)(p-1)+1)^{j-2}}{(j-1)!} \right)^{l_j}$$

where

$$i_{\max} = 1 + \min\left(\left\lfloor \frac{n-1}{p-1} \right\rfloor, k\right)$$

and the set of integers $\{l_2, l_3, \dots, l_{i_{\max}}\}$ satisfies $\sum_{i=2}^{i_{\max}} (i-1)l_i = k$.

PROOF. Since the theorem holds for $k=1$ we assume that it holds for $k-1$ and prove that it holds for k . We start by writing the relation

$$\begin{aligned} & [(p)]_n^k \Big| \prod_{j=1}^{i_{\max}} ((j-1)(p-1)+1)^{l_j} \Big|_n \\ &= \sum_{i=2}^{i_{\max}} \sum_{q=1}^p \sum_{i_1 < i_2 < \dots < i_q} \sum_{\substack{(\lambda_{i_1}, \lambda_{i_2}, \dots, \lambda_{i_q}) \vdash p \\ (\sum_{j=1}^q \lambda_{i_j} = i + p - 2)}} \\ & \times [(p)]_n \cdot \left[\prod_{j=1}^{i_{\max}} ((j-1)(p-1)+1)^{l_j + \lambda_j - \delta_{ji}} \right] \Big| \prod_{j=1}^{i_{\max}} ((j-1)(p-1)+1)^{l_j} \Big|_n \\ & \times [(p)]_n^{k-1} \Big| \prod_{j=1}^{i_{\max}} ((j-1)(p-1)+1)^{l_j + \lambda_j - \delta_{ji}} \Big|_n, \end{aligned} \quad (2)$$

where $(\lambda_{i_1}, \lambda_{i_2}, \dots, \lambda_{i_q}) \vdash p$ means that $\lambda_{i_1}, \lambda_{i_2}, \dots, \lambda_{i_q}$ form a partition of p , i.e. $\lambda_{i_1} \leq \lambda_{i_2} \leq \dots \leq \lambda_{i_q}$ and $\lambda_{i_1} + \lambda_{i_2} + \dots + \lambda_{i_q} = p$. Equation (2) evaluates the coefficient of $\prod_{j=1}^{i_{\max}} ((j-1)(p-1)+1)^{l_j}$ in $[(p)]_n^k$ by summing over all the terms in $[(p)]_n^{k-1}$ that can give rise to that term. The p cycles which the last application of $[(p)]_n$ fuses are grouped into q subsets, with lengths specified by the indices i_1, i_2, \dots, i_q , such that the number of cycles of length $(i_j-1)(p-1)+1$ is λ_{i_j} , i is the index specifying the length, $(i-1)(p-1)+1$, of the cycle inserted as a consequence of the fusion, and the

condition $\sum_{j=1}^q i_j \lambda_{i_j} = i + p - 2$ specifies the fact that the length of that cycle is the sum of the lengths of the p cycles the fusion of which gave rise to it.

To evaluate the first factor on the right-hand side we use Theorem 1. For the second factor we use the induction hypothesis. We consequently obtain

$$\begin{aligned} & [(p)]_n^k \left| \prod_{j=1}^{i_{\max}} ((j-1)(p-1)+1)^{l_j} \right|_n \\ &= k! \prod_{j=2}^{i_{\max}} \left(\frac{((j-1)(p-1)+1)^{j-2}}{(j-1)!} \right)^{l_j} \\ & \times \left\{ \sum_{i=2}^{i_{\max}} \sum_{q=1}^p \sum_{i_1 < i_2 < \dots < i_q} \sum_{\substack{(\lambda_{i_1}, \lambda_{i_2}, \dots, \lambda_{i_q}) \vdash p \\ (\sum_{j=1}^q i_j \lambda_{i_j} = i + p - 2)}} \frac{(p-1)!}{k \prod_{j=1}^q \lambda_{i_j}} ((i-1)(p-1)+1) l_i \right. \\ & \times \left. \prod_{j=2}^{i_{\max}} \left(\frac{((j-1)(p-1)+1)^{j-2}}{(j-1)!} \right)^{l_j} \left(\frac{((i-1)(p-1)+1)^{i-2}}{(i-1)!} \right)^{-1} \right\}. \end{aligned}$$

We shall now show that the expression in curly brackets is equal to unity. This expression can be written in the form

$$\frac{1}{kp} \sum_{i=2}^{i_{\max}} \frac{(i-1)! l_i}{((i-1)(p-1)+1)^{i-3}} \sum_{\substack{i_1, i_2, \dots, i_p \\ (\sum_{j=1}^p i_j = i + p - 2)}} \prod_{j=1}^p \left(\frac{((i_j-1)(p-1)+1)^{i_j-2}}{(i_j-1)!} \right),$$

where the set i_1, i_2, \dots, i_p allows repetitions, so that the sum over this set of indices is equivalent to the sum over the original set i_1, i_2, \dots, i_q as well as over $\lambda_{i_1}, \lambda_{i_2}, \dots, \lambda_{i_q}$, multiplied by $\prod_{j=1}^q \lambda_{i_j}! / p!$. Using the multinomial Abel identity ([13, p. 24])

$$\sum_{\substack{i_1, i_2, \dots, i_p=0 \\ (\sum_{j=1}^p i_j = i)}}^i \prod_{j=1}^p \left(\frac{(i_j(p-1)+1)^{i_j-1}}{i_j!} \right) = \frac{p(i(p-1)+p)^{i-1}}{i!},$$

we write the above expression in the form $(1/k) \sum_{i=2}^{i_{\max}} (i-1) l_i$, which, by equation (1), is equal to unity. \square

Theorem 2 constitutes a generalization of Dénes' result [6], to which it reduces for $p = 2$. The original proof, due to Dénes, is based on establishing a correspondence with a certain enumeration problem involving edges of graphs. It may be instructive to consider the graph-theoretical implications of the present generalization, possibly along the lines discussed by Dénes [6] and, recently, by Moszkowski [12] and by Goulden and Pepper [7], but this is beyond the scope of the present paper.

ACKNOWLEDGEMENTS

This research was supported by the Fund for the Promotion of Research at the Technion. Helpful discussions with Professors D. Chillag, A. Juhász, G. Kalai and R. Meshulam are gratefully acknowledged.

REFERENCES

1. Z. Arad and M. Herzog (eds), *Products of Conjugacy Classes in Groups*, Lecture Notes in Mathematics 1112, Springer-Verlag, Berlin, 1985.
2. E. A. Bertram and V. K. Wei, Decomposing a permutation into two large cycles: an enumeration, *SIAM J. Algebra Discr. Methods*, **1** (1980), 450–461.
3. G. Boccarda, Decompositions d'une permutation d'un ensemble fini en produit de deux cycles, *Discr. Math.*, **23** (1978), 189–205.

4. G. Boccara, Nombre de representations d'une permutation comme produit de deux cycles de longueurs donnees, *Discr. Math.*, **29** (1980), 105–134.
5. G. Boccara, Cycles comme produit de deux permutations de classes donnees, *Discr. Math.*, **38** (1982), 129–142.
6. J. Dénes, The representation of a permutation as the product of a minimal number of transpositions, and its connection with the theory of graphs, *A Matematikai Kutató Intézet Közleményei*, **4** (1959), 63–71.
7. I. P. Goulden and S. Pepper, Labelled trees and factorizations of a cycle into transpositions, *Discr. Math.*, **113** (1993), 263–268.
8. J. Katriel, Products of class operators of the symmetric group, *Int. J. Quantum Chem.*, **35** (1989), 461–470.
9. J. Katriel, A conjecture concerning the evaluation of products of class-sums of the symmetric group, in: *Groups 1993—Galway/St. Andrews*, C. M. Campbell (ed.), London Mathematical Society Lecture Note Series (in press).
10. J. Katriel and J. Paldus, Explicit expression for the product of the class of transpositions with an arbitrary class of the symmetric group, in: *Group Theoretical Methods in Physics*, R. Gilmore (ed.), World Scientific, Singapore, 1987, pp. 503–506.
11. A. Machí, Sur les représentations des permutations impaires, *Europ. J. Combin.*, **13** (1992), 273–277.
12. P. Moszkowski, A solution to a problem of Dénes: a bijection between trees and factorizations of cyclic permutations, *Europ. J. Combin.*, **10** (1989), 13–16.
13. J. Riordan, *Combinatorial Identities*, John Wiley, New York 1968.

Received 29 July 1994 and accepted 21 September 1994

JACOB KATRIEL
Department of Chemistry,
Technion—Israel Institute of Technology,
32000 Haifa, Israel
E mail: chr09k1@technion